

**Lähteet:**

- <sup>1</sup> DAISIE. 7.1.2008 (päivitetty). *Corbicula fluminea* (Dan Minchin toim.). [www.europe-aliens.org/speciesSearch.do](http://www.europe-aliens.org/speciesSearch.do) > Search species: *Corbicula fluminea* > Factsheet. [Viitattu 4.6.2011]
- <sup>2</sup> Karatayev, A.Y., Burkalova, L.E., Padilla, D.K. 2005. Contrasting distribution and impacts of two freshwater exotic suspension feeders, *Dreissena polymorpha* and *Corbicula fluminea*. Kirjassa: R. Dame and S. Olenin (toim.) The Comparative Roles of Suspension Feeders in Ecosystems. NATO Science Series: IV – Earth and Environmental Sciences. Springer, s 239–262.
- <sup>3</sup> Qiu, A., Shi, A. & Komaru, A. 2001. Yellow and brown shell color morphs of *Corbicula fluminea* (Bivalvia: Corbiculidae) from Sichuan Province, China, are triploids and tetraploids. Journal of Shellfish Research 20: 323–328.
- <sup>4</sup> Sousa, R., Freire, R., Rufino, M., Méndez, J., Gaspar, M., Antunes, C. & Guilhermino, L. 2007. Genetic and shell morphological variability of the invasive bivalve *Corbicula fluminea* (Müller 1774) in two Portuguese estuaries. Estuarine, Coastal and Shelf Science 74: 166–174.
- <sup>5</sup> Pathy, D. A. & Mackie, G. L. 1993. Comparative shell morphology of *Dreissena polymorpha*, *Mytilopsis leucophaeata*, and the "quagga" mussel (Bivalvia: Dreissenidae) in North America. Canadian Journal of Zoology, 71: 1012–1023.
- <sup>6</sup> Marelli, D.C. & Gray, S. 1983. Conchological redescrptions of *Mytilopsis sallei* and *Mytilopsis leucophaeata* of the brackish Western Atlantic (Bivalvia: Dreissenidae). Veliger 25: 185–193.
- <sup>7</sup> Verveen, A., Vincx, M. & Degraer, S. 2010. *Mytilopsis leucophaeata* : the brackish water equivalent of *Dreissena polymorpha*? A review. Julk: Van der Velde, G., Rajagopal, S. & bij de Vaate, A. (toim.). The Zebra Mussel in Europe. Leiden, The Neatherlands. Brackhuys Publishers. s. 29–44.
- <sup>8</sup> Orlova, M.I., Telesh, I.V., Berezina, N.A., Anstulevic, A.E., Maximov, A.A. & Litvinchuk, L.F. 2006. Effects of nonindigenous species on diversity and community functioning in the Gulf of Finland (Baltic Sea). Helgol. Mar. Res. 60: 98–105.
- <sup>9</sup> Rosenburg, G. & Ludyanskiy, M.L. 1994. A nomenclatural review of *Dreissena* (Bivalvia: Dreissenidae), with identification of the Quagga Mussel as *Dreissena bugensis*. Can. J. Fish. Aquat. Sci. 51: 1474–1484.
- <sup>10</sup> Therriault, T. W., Docker, M.F., Orlova, M. I., Heath, D.D. & MacIsaac, H.J. 2004. Molecular resolution of the family Dreissenida (Mollusca: Bivalvia) with emphasis on Ponto-Caspian species, including first report of *Mytilopsis leucophaeata* in the Black Sea basin. Molecular phylogenetics and Evolution 30: 479–489.
- <sup>11</sup> van der Velde, G., Rajagopal, S. & bij de Vaate, A. (toim.) 2010. The Zebra Mussel in Europe. Backhuys Publishers. xviii+ 490 ss. ISBN 978-3-8236-1594-1.
- <sup>12</sup> Antsulevich, A.E., Välipakka, P. & Vaittinen, J. 2003. How are the zebra mussels doing in the Gulf of Finland? Proc. Estonian Acad. Sci. Biol. Ecol. 52(3): 268–283.
- <sup>13</sup> Claxton, W.T., Martel, A., Dermott, R.M. & Boulding, E.G. 1997. Discrimination of field-collected juveniles of two introduced dreissenids (*Dreissena polymorpha* and *Dreissena bugensis*) using mitochondrial DNA and shell morphology. Can. J. Fish. Aquat. Sci. 54: 1280–1288.
- <sup>14</sup> Ellis, S. & MacIsaac, H.J. 2009. Salinity tolerance of Great Lakes invaders. Freshwater biology, 54: 77–89.
- <sup>15</sup> Nichols, S.J. & Black, M.G. 1994. Identification of larvae: the zebra mussel (*Dreissena polymorpha*), quagga mussel (*Dreissena rostriformis bugensis*), and Asian clam (*Corbicula fluminea*). Can. J. Zool. 72: 406–417.

- <sup>16</sup> Anonymi 2010. Utah's AIS Management Plan, Appendix A – Dreissenid Mussels: Quagga Mussel *Dreissena bugensis*, Zebra Mussel *Dreissena polymorpha*, False Darkmussel *Mytilopsis leucophaeata*. s. 33–48. Verkossa: [http://wildlife.utah.gov/pdf/AIS\\_plans\\_2010/AIS\\_12iDreissenidMussels-NatalieFinal.pdf](http://wildlife.utah.gov/pdf/AIS_plans_2010/AIS_12iDreissenidMussels-NatalieFinal.pdf) [Viitattu 2.7.2011]
- <sup>17</sup> Dermott, R. & Munawar, M. 1993. Invasion of Lake Erie offshore sediments by *Dreissena*, and its ecological impact. *Can. J. Fish. Aquat. Sci.* 50: 2298–2304.
- <sup>18</sup> Mills, E.L., Rosenberg, G., Spidle, A. P., Ludyanskiy, M., Pligin, Y. & May, B. 1996. A review of the biology and ecology of the quagga mussel (*Dreissena bugensis*), a second species of freshwater Dreissenid introduced to North America. *American Zoologist* 36: 271–286.
- <sup>19</sup> Hayward, P. J. & Ryland, J. S. (toim.) 1995. *Handbook of the Marine fauna of North-West Europe*. Oxford University press, United Kingdom. s. 514–516.
- <sup>20</sup> Fish, J.D. & Fish, S. (2. painos) 1996. *A Student's Guide to the Seashore*. Cambridge University Press.
- <sup>21</sup> Mastitsky, S.E. & Samoilenko, V.M. 2006. The gravel snail, *Lithoglyphus naticoides* (Gastropoda: Hydrobiidae), a new Ponto-Caspian species in Lake Lukomskoe (Belarus). *Aquatic Invasions*, 1(3): 161–170.
- <sup>22</sup> Mastitsky, S.E. 2007. First report of parasites in *Lithoglyphus naticoides* (Gastropoda: Hydrobiidae) from Lake Lukomskoe (Belarus). *Aquatic Invasions*, 2(2): 149–151.
- <sup>23</sup> Fauna Europaea (2010) Fauna Europaea version 2.4. <http://www.faunaeur.org> [Viitattu 31.8.2011]
- <sup>24</sup> Brodin, Y. & Andersson, M.H. 2009. The marine splash midge *Telmatogeton japonicus* (Diptera; Chironomidae) – extreme and alien? *Biol. Invasions*, 11: 1311–1317.
- <sup>25</sup> Raunio, J., Paasivirta, L. & Brodin, Y. 2009. Marine midge *Telmatogeton japonicus* Tokunaga (Diptera: Chironomidae) exploiting brackish water in Finland. *Aquatic invasions*, 4(2): 405–408.
- <sup>26</sup> Tokunaga, M. 1933. Chironomidae from Japan (Diptera) I. Clunioninae. *Philippine Journal of Science* 51: 87-99.
- <sup>27</sup> Paasivirta, L. 2000. Chironomidae (Diptera) of the northern Baltic Sea. Kirjassa: Hoffrichter, O. (toim.) 2000. Late 20th century research on Chironomidae: an anthology from the 13th international symposium on Chironomidae. Freiburg, Germany, September 5–9, 1997. Shaker Verlag, Aachen, Saksa, s. 589–598.
- <sup>28</sup> Lang, W.H. 1980. Balanomorph nauplii of the NW Atlantic shores. *Fisch. Ident. Zooplankton* 163. 6 s.
- <sup>29</sup> Furman, E. 1989. The taxonomic relationships of *Balanus improvisus* (Darwin) populations in the Baltic and the Atlantic. Väitöskirjatutkimus, University College of North Wales. 136 s.
- <sup>30</sup> McLaughlin, P.A. 1980. Comparative morphology of recent crustacea. W.H. Freeman & Company. 177 s.
- <sup>31</sup> Rajasilta, M. & Vuorinen, I. 2008. Suomen murtovesialueen eläinplankton – määrittäysopas. Turun yliopiston, Saaristomeren tutkimuslaitos. 66 s.
- <sup>32</sup> Bradford-Grieve, J.M. 1999. Copepoda, Acartiidae, Acartia, Paracartia, Pteriacartia. Julk.: ICES Identification leaflets for plankton, 181: 1-19. ICES, Copenhagen, Denmark.
- <sup>33</sup> Mauchline, J. 1998. The biology of calanoid copepods. *Crustaceana*, 81(6): 763–764.
- <sup>34</sup> Telesh, I., Postel, L., Heerkloss, R., Mironova, E., Skarlato, S. 2008. Zooplankton of the Open Baltic Sea: Atlas. BMB Publ 20. Meereswiss. Ber. Warnemünde 73:1–251.
- <sup>35</sup> Razouls, C. 1995. Diversité et répartition géographique chez les copépodes pélagiques. 1 - Calanoida. – *Ann. Inst. océanogr. (Paris)* 71: 81–401.

- <sup>36</sup> Leppäkoski, E. & Olenin, S. 2000. Non-native species and rates of spread: lessons from the brackish Baltic Sea. – *Biological Invasions* 2: 151–163.
- <sup>37</sup> Silina, N.I. [Силина, Н.И.] 1989. Виды рода *Acartia* в планктоне восточной части Финского Залива Балтийского моря. – *Труды Зоол. ин-та АН СССР* [Species of the genus *Acartia* in the plankton of the eastern part of the Gulf of Finland, the Baltic Sea. (USSR Acad. Sci., Proc. Zool. Inst., Leningrad) 205: 108–118].
- <sup>38</sup> Aquatic Nuisance Species Research Program, lokakuu 2006 (päivitetty). Fishhook waterflea - *Cercopagis pengoi*. <http://el.erdc.usace.army.mil/ansrp/ansrp.html> > Species of concern > Review existing profiles > Species profiles > *Cercopagis pengoi* - Fishhook Waterflea. toim.: Crosier, D.M., Molloy, D.P. & Grigorovich, A. [Viitattu 23.12.2011]
- <sup>39</sup> Rivier, I.K. 1998. The predatory Cladocera (Onychopoda: Podonidae, Polyphemidae, Cercopagidae) and Leptodorida of the world. Backhuys Publishers, Leiden, 213 s.
- <sup>40</sup> Martin, J.W. & Cash-Clark, C.E. 1995. The external morphology of the onychopod 'cladoceran' genus *Bythotrephes* (Crustacea, Branchiopoda, Onychopoda, Cercopagididae), with notes on the morphology and phylogeny of the order Onychopoda. *Zoologica Scripta*, 24(1): 61–90.
- <sup>41</sup> Grigorovich, I.A., MacIsaac, H.J., Rivier, I.K., Aladin, N.V. & Panov, V.E. 2000. Comparative biology of the predatory cladoceran *Cercopagis pengoi* from Lake Ontario, Baltic Sea and Caspian Sea. *Arch. Hydrobiol.*, 149: 23–50.
- <sup>42</sup> Simm, M. & Ojaveer, H. 2006. Taxonomic status and reproduction dynamics of the non-indigenous *Cercopagis* in the Gulf of Riga (Baltic Sea). *Hydrobiologia*, 554: 147–154.
- <sup>43</sup> Eloranta, P. 2001. Opas Suomen sisävesien eläinplanktonin tuntemukseen (Rataseläimet, vesikirput, hankajalkaiset). *Limnologian ja ympäristönsuojelun laitos/limnologia*. Helsingin Yliopisto.
- <sup>44</sup> Panov, V., Rodionova, N., Bolshagin, P. & Bychek, E. 2007. Invasion biology of Ponto-Caspian onychopod cladocerans (Crustacea: Cladocera: Onychopoda). *Hydrobiologia*, 590: 3–14.
- <sup>45</sup> Särkkä, J. 1984. Sisävesien eläinplanktonituntemuksen alkeita. Jyväskylän yliopiston biologian laitoksen opetusmonisteita. 25 s.
- <sup>46</sup> Rodionova, N., Krylov, P. & Panov, V. 2005. Invasion of the Ponto–Caspian Predatory Cladoceran *Cornigerius maeoticus maeoticus* (Pengo, 1879) into the Baltic Sea. *Oceanology* 45: 66–68.
- <sup>47</sup> Australian museum 2009. Crustacea.net – An information retrieval system for crustaceans of the world. [www.crustacea.net/](http://www.crustacea.net/). [Viitattu 13.9.2011]
- <sup>48</sup> Anderson, G. 20.1.2010. Cumacea Classification. <http://peracarida.usm.edu/> > Cumacea. [Viitattu 28.8.2011]
- <sup>49</sup> Bacescu, M. 1969. Otryad kumovye – Cumacea. Julk.: F. D. Mordukhai-Boltovskoi (toim.) *Opredelitel fauny Chernogo i Azovskogo morei*, T. 2. Svobodnozhivuschie bespozvonochnye. Rakoobraznye, s. 381–402. [Order Cumacea. Julk.: A Key of the Black Sea and Sea of Azov fauna. V. 2. Free-living invertebrates. Crustaceans.]
- <sup>50</sup> Lomakina, N.B. 1958. Kumovye raki (Cumacea) morei SSSR. Moscow, Leningrad, AN SSSR, 301 s. (Opredeliteli po faune SSSR, izdavaemye Zoologicheskim Institutom AN SSSR, 66) [Cumacea of the USSR seas].
- <sup>51</sup> Antsulevich, A.E. 2005. First finding of Cumacea crustaceans in the Gulf of Finland.- *Vestnik Sankt-Peterburgskogo Universiteta, Seria 3 (Biologija 1)*: 84-87.
- <sup>52</sup> Kusakin, O.G. 1969. Otryad ravnogie – Isopoda. Julk.: F. D. Mordukhai-Boltovskoi (toim.) *Opredelitel fauny Chernogo i Azovskogo morei*, T. 2. Svobodnozhivuschie bespozvonochnye. Rakoobraznye. S. 408–440. [Order Isopoda. Julk.: A Key of the Black Sea and Sea of Azov fauna. V. 2. Free-living invertebrates. Crustaceans.]
- <sup>53</sup> Kusakin, O.G. 1988. Morskie i solonovatovodnye ravnogie rakoobraznye (Isopoda) kholodnyh i umerennykh vod severnogo polushariya. T. III. Podotryad Asellota. Ch. I. Semeistva

Janiridae, Santidae, Dendrotionidae, Munnidae, Paramunnidae, Haplomunnidae, Mesosignidae, Haploniscidae, Mictosomatidae, Ischnomesidae. Leningrad, Nauka, 502 s. (Opredeliteli po faune SSSR, izdavaemye Zoologicheskim Institutom AH SSSR, 152). [Marine and brackish-water isopods of cold and moderate waters of the North Hemisphere. V. 3. Suborder Asellota. Part 1. Families Janiridae, Santidae, Dendrotionidae, Munnidae, Paramunnidae, Haplomunnidae, Mesosignidae, Haploniscidae, Mictosomatidae, Ischnomesidae.]

- <sup>54</sup> Wilson, G.D.F. & Wagele, J.W. 1994. A systematic review of the family Janiridae (Crustacea, Isopoda, Asellota). *Invertebrate Taxonomy*, 8(3): 683–747.
- <sup>55</sup> Orlova, M.I., Telesh, I.V., Berezina, N.A., Antsulevich, A.E., Maximov, A.A. & Litvinchuk, L.F. 2006. Effects of nonindigenous species on diversity and community functioning in the eastern Gulf of Finland (Baltic Sea). *Helgol. Mar. Res.*, 60: 98-105.
- <sup>56</sup> Tobias, W., Wegmann, A. & Bernerth, H. 2005: Jaera istri oder Jaera sarsi? - Zum taxonomischen Status der "Donauassel" (Isopoda, Asellota: Janiridae). - *Umwelt und Geologie: Faunistisch-ökologische Untersuchungen des Forschungsinstitutes Senckenberg im hessischen Main: 5-14, Wiesbaden*.
- <sup>57</sup> Bousfield, E. L. 1973: *Shallow-water gammaridean Amphipoda of New England*. — Comstock Publ. Assoc., div. Cornell Univ. Press, Ithaca, 312 pp.
- <sup>58</sup> Silfverberg, H. 1999. A provisional list of Finnish Crustacea. *Memoranda Soc. Fauna Flora Fennica* 75: 15–37
- <sup>59</sup> Hurley, D.E. 1958. A key to the Families of New Zealand Amphipods. *Tuatara* 7(2): 74.
- <sup>60</sup> Konopacka, A. & Jazdzewski, K. 2002. *Obesogammarus crassus* (G.O. Sars, 1894) - one more Ponto-Caspian gammarid species in Polish waters. *Frag. faun.* 45: 19–26.
- <sup>61</sup> Bousfield, E.L. 1858. Fresh-water amphipod crustaceans of glaciated North America. *The Canadian Field-Naturalist* 72(2): 55–113.
- <sup>62</sup> Aquatic Nuisance Species Research Program, lokakuu 2006 (päivitetty). *Echinogammarus ischnus* – Amphipod. <http://el.erdc.usace.army.mil/ansrp/ansrp.html> > Species of concern > Review existing profiles > Species profiles > *Echinogammarus ischnus* – Amphipod. toim.: Crosier, D.M., Molloy, D.P. & Grigorovich, A. [Viitattu 23.12.2011]
- <sup>63</sup> Mordukhai-Boltovskoi, F. D., Greze, I. I. & Vassilenko, C. B. 1969. Otryad amfipody, ili raznogie – Amphipoda. *Julk.: F. D. Mordukhai-Boltovskoi (toim.) Opredelitel fauny Chernogo i Azovskogo morei, T. 2. Svobodnozhivuschie bespozvonochnye. Rakoobraznye*, s. 440–524. [Order Amphipoda. In: *A Key of the Black Sea and Sea of Azov fauna. V. 2. Free-living invertebrates. Crustacens.*]
- <sup>64</sup> Arbačiauskas, K. 2005. The distribution and local dispersal of Ponto-Caspian peracarida in Lithuanian fresh waters with notes on *Pontogammarus robustoides* population establishment, abundance and impact. *Oceanological and Hydrobiological Studies*, Vol. XXXIV (Supplement 1): 93–111.
- <sup>65</sup> Karaman, G.S. & Barnard, J.L. 1979. Classificatory revisions in Gammaridean Amphipoda (Crustacea), part 1. *Proceedings of the Biological Society of Washington*, 92 (1): 106–165.
- <sup>66</sup> Lincoln, R.J. 1979. *British Marine Amphipoda: Gammaridea*. British Museum (Natural History), London. 658 s.
- <sup>67</sup> Spicer, J.I. & Janas, U. 2006. The beachflea *Platorchestia platensis* (Krøyer, 1845): a new addition to the Polish fauna (with a key to Baltic talitrid amphipods). *Oceanologia* 48(2): 287–295.
- <sup>68</sup> Persson, L.-E. 2001. Dispersal of *Platorchestia platensis* (Krøyer) (Amphipoda: Talitridae) along Swedish coasts: A slow but successful Process. *Estuarine, Coastal and Shelf Science* 52: 201–210.
- <sup>69</sup> Dekker, W. 1978. Strandvlooiën (Talitridae). *Tabellenserie van de Strandwerkgemeenschap*. 24 s.

- <sup>70</sup> Akbulut, M. & Sezgin, M. 2000. *Orchestia cavimana* Heller, 1865 (Amphipoda, Talitridae) in the Turkish Black Sea Fauna. Turkish J. Marina Sciences 6: 241–244.
- <sup>71</sup> Herkül, K., Kotta, J. & Ilmar, K. 2006. Distribution and population characteristics of the alien talitrid amphipod *Orchestia cavimana* in relation to environmental conditions in the Northeastern Baltic Sea. Helgol. Mar. Res. 60: 121–126.
- <sup>72</sup> Bazikalova, A. Y. 1945. Amfipody ozera Baikal. Trudy Baikalskoi Limnologicheskoi Stantsii AN SSSR, 11. 440 s.
- <sup>73</sup> Stebbing, T.R.R. 1899. Amphipoda from the Copenhagen Museum and other sources. Part 2. Transaction of the Linnean Society of London, 7(8): 395–432.
- <sup>74</sup> Berezina, N.A., Khlebovich, V.V., Panov, V.E. & Zaporozhets, N.V. 2001. Salinity tolerance of the amphipod *Gmelinoides fasciatus* (Stebb) introduced into the Gulf of Finland basin (the Baltic Sea). Doklady Biological Sciences, 379: 366–368.
- <sup>75</sup> Berezina, N.A. & Panov, V.E. 2004. Distribution, population structure and salinity tolerance of the invasive amphipod *Gmelinoides fasciatus* (Stebbing) in the Neva Estuary (Gulf of Finland, Baltic Sea). Hydrobiologia 514: 199–206.
- <sup>76</sup> Uryupova, E.F. 2008 Morphologicheskii, phylogeneticheskii i ekologicheskii obzor ponto-kaspiiskikh korofiid (Corophiinae, Corophiidae, Amphipoda). Avtoreferat na soiskanie uchenoi stepeni kandidata biologicheskikh nauk. Moskva, 25 s. [Morphological, phylogenetic and ecological review of Ponto-Caspian corophiids (Corophiinae, Corophiidae, Amphipoda). Väitöskirja abstrakti]
- <sup>77</sup> Köhn, J. 1992. Mysidacea of the Baltic Sea – State of the art. Julk.: Köhn, J., Jones, M.B. & Moffat, A. (toim.). Taxonomy, Biology and Ecology of (Baltic) mysids (Mysidacea: Crustacea). International Expert Conference, September 1991, Hiddensee, Germany, Rostock University, Germany . s. 5–23.
- <sup>78</sup> Mauchline, J. 1980. The Biology of Mysids – The species of mysids and key to genera. Julk.: Russel, F.S. & Yonge, M. (toim.). Advances in Marine Biology, vol 18. Academic Press Inc (London) Ltd. s. 6–39.
- <sup>79</sup> Tattersall, W.M. & Tattersall, O.L. 1951. The British Mysidacea. The Ray Society, London. 460 s.
- <sup>80</sup> Kelleher, B., van der Velde, G., Wittman, K.J., Faasse, M.A. & bij de Vaate, A. 1999. Current status of the freshwater mysidae in the Neatherlands, with records of *Limnomysis benedeni* Czerniavskyi, 1882, a Ponto-Caspian species in Dutch Rhine branches. Bulletin Zöologische Museum, Vol. 16, No. 13. Universiteit van Amsterdam.
- <sup>81</sup> Salemaa, H. & Hietalahti, V. 1993. *Hemimysis anomala* G.O. Sars (Crustacea: Mysidacea) – Immigration of a Pontocaspian mysid into the Baltic Sea. Ann. Zool. Fennici 30: 271–276.
- <sup>82</sup> Băcescu, M. 1954. Fauna Republicii Populare Romine. Crustacea. Mysidacea. IV(3): 1–126. – Acad. Rep. Pop. Romine.
- <sup>83</sup> Pothoven, S.A., Grigorovich, I.A., Fahnenstiel, G.L. & Balcer, M.D. 2007. Introduction of the Ponto-Caspian bloody-red mysid *Hemimysis anomala* into the Lake Michigan basin. Journal of Great Lakes Research 33(1): 285–292.
- <sup>84</sup> Herkül, K., Kotta, J., Püss, T. & Kotta, I. 2009. Crustacean invasions in the Estonian coastal sea. Estonian Journal of Ecology, 58(4): 313–323.
- <sup>85</sup> Daneliya, M. & Petryashov, V. 2011. Rediscription of three species and a subspecies of the mysid genus *Paramysis* (Mysida, Mysidae) from the ponto-Caspian basin. Crustaceana 84(7): 797–829.
- <sup>86</sup> Ovčarenko, I., Audzijonytė, A. & Gasiūnaitė, A.R. 2006. Tolerance of *Paramysis lacustris* and *Limnomysis benedeni* (Crustacea, Mysida) to sudden salinity changes: implications for ballast water treatment. Oceanologia, 48(S): 231–242.

- <sup>87</sup> Fish, J.D. & Fish, S. (2. ed.) 1996. A Student's Guide to the Seashore. Cambridge University Press. 584 s.
- <sup>88</sup> Janas, U., Zarzycki, T. & Kozik, P. 2004. *Palaemon elegans* – a new component of the Gulf of Gdańsk macrofauna. *Oceanologia* 46(1): 143–146.
- <sup>89</sup> Barnes, R.S.K. (1994). The Brackish-Water fauna of Northwestern Europe: a guide to brackish-water habitats, ecology, and macrofauna for field workers, naturalists, and students. Cambridge University Press. 287 s.
- <sup>90</sup> The McGraw-Hill Companies. Access Science. Encyclopedia article: "Decapoda, Crustacea". <http://accessscience.com/overflow.aspx?searchStr=Decapoda+%28Crustacea%29&stype=10&term=Decapoda+%28Crustacea%29&rootID=792007> [Viitattu 29.6.2011]
- <sup>91</sup> Davie, P.J.F. & Ng, N.K. 2007. Two new subfamilies of Varunidae (Crustacea; Brachyura), with description of two new genera. *The Raffles Bulletin of Zoology, Supplement* 16: 257–272.
- <sup>92</sup> Martin, J. W. & Davis, G. E. 2001. An Updated Classification of the Recent Crustacean. Natural history Museum of Los Angeles County, Science Series 39. Los Angeles, United States of America. ISSN 1-891276-27-1.
- <sup>93</sup> Schubart, C. D., Neigel, J.E. & Felder D. L. 2000. Molecular phylogeny of mud crabs (Brachyura: Panopeidae) from the northwestern Atlantic and the role of morphological stasis and convergence. *Marine Biology* 137: 11–18.
- <sup>94</sup> Rahtbun, M. J. 1930. The Cancroid crabs of America of the families Euryalidae, Portunidae, Atelecyclidae, Cancridae and Xanthidae. Smithsonian Institution, United States National Museum, Bulletin 152. Washington, United States. s. 455–458.
- <sup>95</sup> Marine Species Identification Portal. 2011 (päivityvä). Chinese mitten crab (*Eriocheir sinensis*) – Crabs of Japan. <http://species-identification.org/> > Search for species "Chinese mitten crab". [Viitattu 20.4.2011.]
- <sup>96</sup> Hamr P (2002) *Orconectes*. In: Holdich DM (ed) Biology of freshwater crayfish. Blackwell Science, Oxford, pp 585-608
- <sup>97</sup> Benejam, L., Saura-Mas, S. & Saperas, A. 2011. First record of the spiny-cheek crayfish *Orconectes limosus* (Rafinesque, 1817) introduced to the Iberian Peninsula. *Aquatic Invasions* 6(S1): S111–S113.
- <sup>98</sup> Jaszczolt, J. & Szaniawska, A. 2011. The spiny-cheek crayfish *Orconectes limosus* (Rafinesque, 1817) as an inhabitant of the Baltic Sea – experimental evidence for its invasion of brackish waters. *Int J Ocean Hydrob.* 40(3):52–60.
- <sup>99</sup> Pârvulescu 2010. Crayfish field guide of Romania. Editura Bioflux, Cluj-Napoca. [http://www.editura.bioflux.com.ro/docs/Crayfish\\_guide.pdf](http://www.editura.bioflux.com.ro/docs/Crayfish_guide.pdf)
- <sup>100</sup> Grabowski, M., Jazdzewski, K. & Konopacka, A. 2005. Alien Crustacea in Polish waters – Introduction and Decapoda. *Oceanological and Hydrobiological Studies* XXXIV(S1): 43-61.
- <sup>101</sup> Holdich, D.M., Reynolds, J.D., Souty-Grosset, C., & Sibley, P.J. 2009. A review of the ever increasing threat to European crayfish from non-indigenous crayfish species. *Knowledge and Management of Aquatic Ecosystems* (2009): 394-395, 11.
- <sup>102</sup> Anon 2010. *Orconetes limosus* Amerikansk dvärgkräfta. Främmande arter i Svenska hav faktablad. [http://www.frammandearter.se/5arter/pdf/Orconectes\\_limosus.pdf](http://www.frammandearter.se/5arter/pdf/Orconectes_limosus.pdf). [Viitattu 23.2.2012].
- <sup>103</sup> Gorokhova, E. & Lehtiniemi, M. 2010. Considering evidence for *Mnemiopsis* invasion in European waters. *Journal of plankton research* 32 (1): 93-95.
- <sup>104</sup> Kube, S. & Postel, L. (julkaisematon). Life history stages of *Mnemiopsis leidyi*. Julkaisematon moniste, Baltic Sea Research Institute. 4 s.
- <sup>105</sup> Mayer, A. G. 1912. Ctenophores of the Atlantic Coast of North America. Carnegie Institution of Washington, julkaisu nro 162. 56 s.

- <sup>106</sup> Bryant, J.A. 2002. Order Charales. The freshwater algal flora of the British isles: an identification guide to freshwater and terrestrial algae. 593-612.
- <sup>107</sup> Tolstoy, A. & Österlund, K. 2003. Alger vid Sveriges Östersjökust - en fotoflora. ArtDatabanken, Uppsala. 282s.
- <sup>108</sup> Torn, K., Martin, G., Kukkk, H. & Trei, T. 2004. Distribution of charophyte species in Estonian coastal water (NE Baltic Sea). *Scientia Marina* 68, 129-136.
- <sup>109</sup> Appelgren, K., Snickars, M. & Mattila, J. 2004. *Chara connivens* Saltzm.Ex. A. Braun 1835 found in the Åland archipelago - a new species to Finland. *Memoranda Societas Fauna Flora Fennica* 80, 11-13.
- <sup>110</sup> Ljungberg, R., Pikkarainen, A., Lehtiniemi, M. & Urho, L. 2011. Vieraslajien havaitseminen Suomen merialueen seurannoissa. *Suomen ympäristö* 10, 68s.
- <sup>111</sup> Puntila, R.I. 2007. Basinventering av potentiellt viktiga *Chara*-vikar på norra Åland. *Forskningsrapporter från Husö biologiska station*, 68s.
- <sup>112</sup> Martin, G., Torn, K. & Kotta, J. 2003. Recent changes in distribution pattern of charophyte species *Chara connivens* in Estonian coastal waters of the Baltic Sea. Abstracts of the Third International Conference of Marine Bioinvasions.
- <sup>113</sup> Josefsson, M. 2011. NOBANIS - Invasive Alien Species Fact Sheet *Elodea canadensis*, *Elodea nuttallii* and *Elodea callitrichoides*. Online Database of the European Network on Invasive Alien Species – NOBANIS at <<http://www.nobanis.org/files/factsheets/Elodea.pdf>>
- <sup>114</sup> Främmande arter i Svenska Hav 2006. Canadian waterweed (*Elodea canadensis*) och Nuttall's waterweed (*Elodea nuttallii*). Fact sheet.
- <sup>115</sup> Sarvala, J. 2005. Littoistenjärven ekologisen tilan kehitys ja hoitovaihtoehdot.
- <sup>116</sup> Laita, M., Tarvainen, A., Mäkinen, A., Sammalkorpi, I., Kemppainen, E., Laitinen, L. 2007. Uposkasvien runsastumisesta 2000-luvun alussa. *Suomen ympäristö* 20/2007, Helsinki.
- <sup>117</sup> Lehtonen, J. 2000. Vesirutto ja karvalehti täyttävät ajoittain Littoistenjärven. *Aurora* 1/2000.
- <sup>118</sup> Hintikka, T.J. 1917. Kanadalaisesta vesirutosta ja sen levenemisestä Euroopassa, eritoten Suomessa. *Luonnon Ystävä* 21, 79-90.
- <sup>119</sup> Främmande arter i Svenska Hav 2011. *Elodea canadensis* Vattenpest. Fact sheet [http://www.frammandearter.se/5arter/pdf/Elodea\\_canad\\_nuttall.pdf](http://www.frammandearter.se/5arter/pdf/Elodea_canad_nuttall.pdf).
- <sup>120</sup> Främmande arter i Svenska Hav 2011 *Elodea nuttallii* Smal vattenpest. Fact sheet [http://www.frammandearter.se/5arter/pdf/Elodea\\_canad\\_nuttall.pdf](http://www.frammandearter.se/5arter/pdf/Elodea_canad_nuttall.pdf).

**Lähteitä polttiais- ja sammaleläinten tunnistukseen:**

- Hyman, L.H. 1940. The invertebrates. Vol. 1, Protozoa through Ctenophora. American Museum of Natural History, New York. 726 s.
- Hyman, L.H. 1959. The invertebrates. Vol. 5, Smaller coelomate groups : Chaetognatha, Hemichordata, Pogonophora, Phoronida, Ectoprocta, Brachiopoda, Sipunculida : The coelomate Bilateria. American Museum of Natural History, New York. 783 s.
- Kramp, P.L. 1959. The hydromedusae of the Atlantic Ocean and adjacent waters. Dana – report No. 46. Copenhagen.
- Mayer, A.G. 1910. Medusae of the World. Carnegie Institution of Washington publication, No. 109. Washington.
- Schubert, P. 2004. Revision of the European athecate hydroids and their medusa (Hydrozoa, Cnidaria): families Oceanidae and Pachycordylidae. *Revue Suisse de Zoologie* 111(2): 315–369.